bioRxiv preprint doi: https://doi.org/10.1101/2022.10.13.512134; this version posted October 14, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

Role of spike in the pathogenic and antigenic behavior of SARS-CoV-2 BA.1 Omicron

Da-Yuan Chen[1,2], Devin Kenney[2,3], Chue Vin Chin[1,2], Alexander H. Tavares[1,2], Nazimuddin Khan[1,2], Hasahn L. Conway[1,2], GuanQun Liu[4], Manish C. Choudhary[5,6], Hans P. Gertje[2], Aoife K. O'Connell[2], Darrell N. Kotton[7,8], Alexandra Herrmann[9], Armin Ensser[9], John H. Connor[2,3], Markus Bosmann[10,11,12], Jonathan Z. Li[5,6], Michaela U. Gack[4], Susan C. Baker[13], Robert N. Kirchdoerfer[14], Yachana Kataria[11], Nicholas A. Crossland[2,11], Florian Douam[2,3], Mohsan Saeed[1,2#]

1. Department of Biochemistry, Boston University School of Medicine, Boston, MA, USA
2. National Emerging Infectious Diseases Laboratories, Boston University, Boston, MA, USA
3. Department of Microbiology, Boston University School of Medicine, Boston, MA, USA
4. Cleveland Clinic Florida Research and Innovation Center, Port St. Lucie, FL, USA
5. Brigham and Women's Hospital, Boston, MA, USA
6. Harvard Medical School, Cambridge, MA, USA
7. Center for Regenerative Medicine of Boston University and Boston Medical Center, Boston, MA, USA
8. The Pulmonary Center and Department of Medicine, Boston University School of Medicine, Boston, MA, USA